



Contribution ID: 347

Type: Oral Presentation

Search for a new spin-0 scalar and a spin-1 boson using Run2 ATLAS detector data

Thursday 10 July 2025 11:30 (20 minutes)

We present a search for a spin-1 boson together with a spin-0 scalar where the additional scalar decays into a four lepton final state ($\ell = \mu$ or e) via two intermediate dark vector bosons in the following decay channel $S \rightarrow Z_d Z_d \rightarrow 4\ell$. In this scenario, the targeted additional scalar (S) mass ranges is between 20 GeV and 1 TeV where we exclude the Higgs boson mass window of $115 \text{ GeV} < m_S < 130 \text{ GeV}$ while the dark vector boson has a mass between 15 and 300 GeV. The search is conducted using $p-p$ collision data collected using the ATLAS detector at the LHC which corresponds to a center of mass energy of $\sqrt{S} = 13 \text{ TeV}$ and an integrated luminosity of 139 fb^{-1} . There were no significant excesses observed. Therefore, a 95% upper limit was set on the cross-section $\times$ branching ratio as a function of the mass of both particles m_S and m_{Z_d} .

Apply for student award at which level:

PhD

Consent on use of personal information: Abstract Submission

Yes, I ACCEPT

Primary author: MAPEKULA, Xola (University of Johannesburg)

Presenter: MAPEKULA, Xola (University of Johannesburg)

Session Classification: Nuclear, Particle and Radiation Physics-2

Track Classification: Track B - Nuclear, Particle and Radiation Physics